

S.E - CIVIL - SEM-III CBCS

(3hours)

[Total marks: 80]

14 NOV 2019



- N.B. 1) Question No. 1 is compulsory.
 2) Answer any Three from remaining
 3) Figures to the right indicate full marks

1. a) Find Laplace transform of $f(t) = e^{-4t} \sin 3t \cos 2t$. 5b) Show that the set of functions $f(x) = 1, g(x) = x$ are orthogonal on $(-1,1)$.
 Determine the constants a and b such that the function $h(x) = -1 + ax + bx^2$ is orthogonal to both $f(x)$ and $g(x)$. 5c) Evaluate $\int_C (z^2 - 2\bar{z} + 1) dz$ where C is the circle $|z| = 1$. 5d) Compute the Spearman's Rank correlation coefficient R and Karl Pearson's correlation coefficient r from the following data, 5

x	12	17	22	27	32
y	113	119	117	115	121

2. a) Using Laplace transform, evaluate $\int_0^\infty e^{-t} \int_0^t \frac{\sin u}{u} du dt$. 6b) Find an analytic function $f(z) = u + iv$, if 6
 $u = e^{-x} \{(x^2 - y^2) \cos y + 2xy \sin y\}$.c) Obtain Fourier series of $f(x) = x^2$ in $(0, 2\pi)$. Hence, deduce that 8

$$\frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$$

3. a) Using Bender-Schmidt method, solve $\frac{\partial^2 u}{\partial x^2} - \frac{\partial u}{\partial t} = 0$, subject to the conditions, $u(0, t) = 0, u(4, t) = 0, u(x, 0) = x^2(16 - x^2)$ taking $h = 1$, for 3 minutes. 6b) Using convolution theorem, find the inverse Laplace transform of 6

$$F(s) = \frac{s^2 + s}{(s^2 + 1)(s^2 + 2s + 2)}$$

c) Using Residue theorem, evaluate

$$\text{i) } \int_0^{2\pi} \frac{d\theta}{2 + \cos \theta} \quad \text{ii) } \int_C \frac{z^2}{(z+1)^2(z-2)} dz, \quad C: |z| = 1.5 \quad \text{8}$$

4. a) Solve by Crank –Nicholson simplified formula $\frac{\partial^2 u}{\partial x^2} - 16 \frac{\partial u}{\partial t} = 0$,
 $u(0, t) = 0, u(1, t) = 200t, u(x, 0) = 0$ taking $h = 0.25$ for one-time step.

b) Obtain the Laurent series which represent the function

$$f(z) = \frac{4z+3}{z(z-3)(z+2)} \text{ in the regions, i) } 2 < |z| < 3 \quad \text{ii) } |z| > 3$$

- c) Solve $(D^2 - 3D + 2)y = 4e^{2t}$ with $y(0) = -3$ and $y'(0) = 5$ where $D \equiv \frac{d}{dt}$

5. a) Find the bilinear transformation under which $1, i, -1$ from the z -plane are mapped onto $0, 1, \infty$ of w -plane.

b) Find the Laplace transform of

$$f(t) = \begin{cases} t, & 0 < t < \pi \\ \pi - t, & \pi < t < 2\pi \end{cases} \text{ and } f(t + 2\pi) = f(t).$$

- c) Obtain half range Fourier cosine series of $f(x) = x, 0 < x < 2$. Using Parseval's identity, deduce that –

$$\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$$

6. a) Using contour integration, evaluate:

$$\int_{-\infty}^{\infty} \frac{x^2 + x + 2}{x^4 + 10x^2 + 9} dx$$

- b) Using least square method, fit a parabola, $y = a + bx + cx^2$ to the following data,

x	-2	-1	0	1	2
y	-3.150	-1.390	0.620	2.880	5.378

- c) Determine the solution of one-dimensional heat equation $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$ under the boundary conditions $u(0, t) = 0, u(l, t) = 0, u(x, 0) = x, (0 < x < l), l$ being the length of the rod.