Q.P. Code:05601

[Time: 21/2Hours]

[ Marks:75]

Please check whether you have got the right question paper.

N.B:

- 1. All questions are compulsory.
- 2. Figures to the right indicate marks.





- a) Choose the best choice for the following questions:
  - 1) Let f be defined on an interval, and let x1 and x2 be points on the interval, then f is said to be a
    - p)  $f(x_1)=f(x_2)=0$
    - q)  $f(x_1)=f(x_2)=1$
    - r)  $f(x_1)=f(x_2)=k$
    - s) all of these
  - 2) if f''(a) exists and f has an inflection point at x = a, then
    - p) f''(a) > 0
    - q) f''(a) < 0
    - r) f''(a)=0
    - s) none of these
  - 3) If a function f is continuous on an interval [a,b], then which of the following is true:
    - p) f is integrable on [a,b]
    - q) f is differentiable on [a,b]
    - r) Either (P) or (q)
    - s) None of these
  - 4) the graph of a function of two variables is a surface in
    - p) 1-space
    - q) 2-space
    - r) 3-space
    - s) None of theses
  - 5) which of the following is true about the function  $f(x,y) = \frac{xy}{1+x^2+y^2}$ ?
    - p) Continuous everywhere
    - q) Continuous except where 1+x2+y2=0
    - r) Either (p) or (q)
    - s) Neither (p) nor (q)
- b) Fill in the blanks for the following questions:
  - 1. Two non-negative numbers, x and y, have a sum equal to 10. The largest possible product of the two numbers is obtained by maximizing f(x) = ----- for x in the interval.
  - 2. If y = f(x) is a smooth curve on the interval [a, b] then the arc length of this curve over [a, b] is defined
  - 3. A solution of a differential equation  $\frac{dy}{dx}$  y=0 is given by -----
  - 4. If f (x, y)=  $\sqrt{y+1}\log(x^2-y)$ , the value of f (e,0) is given by------
  - 5. The value of  $\lim (x, y) \rightarrow (3,2)x\cos(\pi y) = \cdots$

(Turn Over)

05

## Q.P. Code:05601


- c) State true or false for the following questions:
  - 1. Newtons Method uses the tangent line to y=f(x) at  $x=x_n$  to compute  $x_n+1$ .

  - The differential equation \$\frac{d^2y}{dx^2} = \frac{dy}{dx}\$ has a solution which is constant.
    If f and g are functions of two variables such that f + g and f g are both continuous, then f and g are themselves continuous.
  - 4. If  $f(x,y) \rightarrow L$  as  $(x,y) \rightarrow (x_0,y_0)$ , then  $f(x,y) \rightarrow L$  as  $(x,y) \rightarrow (x_0,y_0)$  along any smooth curve.
  - 5. A function f of two variables is said to have an absolute maximum at a point  $(x_0, y_0)$  if  $f(x_0, y_0) \le f(x_0, y_0)$ y) for all points (x, y) in the domain of f.
- ANSWER ANY three of the following question: Q.2
  - a) Find the intervals on which  $f(x)=x^2-4x+3$  is increasing and the intervals on which it is decreasing.
  - b) Use first and second derivative tests to show that  $f(x)=x^3-3x+3$  has a relative minimum at x=1 and a relative maximum at x=-1.
  - c) Locate the critical points of  $f(x) = 3x^4 + 12x$ .
  - d) Find the absolute maximum and minimum values of  $f(x) = 4x^2 12x + 10$  in [1,2].
  - e) A firm determines that x units of its product can be sold daily at p Rupees per unit, where x=1000p. The cost of producing x units per day is C(x) = 3000 + 20x.
    - 1. Find the revenue function R(x)
    - 2. Find the profit function P(x)
    - 3. Assuming that the production capacity is at most 500 units per day, determine how many units the company must produce and sell each day to maximize the profit
    - 4. Find the maximum profit.
  - f) The equation x<sup>3</sup>-x-1=0 has one real solution. Approximate it by Newtons Method.
- Q.3 Answer any THREE of the following questions:
  - a) Find the area under the curve  $y=3\sqrt{x}$  over the interval [1,4].
  - b) Find the area of the region enclosed by  $x=y^2$  and y=x-2, integrating with respect to y.
  - c) Find the approximate value of  $\int_{1}^{2} \frac{1}{r} dx$  using Simpson's rule with n=14.
  - d) Solve differential equation  $\frac{dy}{dx} = 2(1 + y^2)x$ .
  - e) Use Euler's Method with a step size of 0.5 to find approximate solution of the initial-value problem  $\frac{dy}{dx} = y^{\frac{1}{3}}, y(x) = 1$  over  $0 \le x \le 4$ .
  - Solve the differential equation  $\frac{dy}{dx} y = e^x$  by the method of integrating factors.

(Turn Over)

15

15






15


- Q.4 Answer any THREE of the following question:
  - a) Find  $\lim_{(x,y)\to(0,0)} \frac{xy}{x^2+y^2}$  (1) along x-axis and (2) along the parabola  $y=x^2$ .
  - b) Determine whether the limit exists. If so, find its value.

$$\lim_{(x, y) \to (0,0)} \frac{1-x^2-y^2}{x^2+y^2}$$

c) Find  $f_x(2,1)$  and  $f_y(1,2)$  for the function  $f(x,y)=10x^2y^4-6xy^2+10x^2$ .



- e) Find parametric equations of the tangent line to the curve of intersection of the paraboloid  $z=x^2+y^2$ And the ellipsoid  $3x^2+2y^2+z^2=9$  at the point (1,1,2).
- f) Find all relative extrema and saddle points of f (x,y)=1- $x^2-y^2$ .



- a) Let  $f(x)=x^2+px+q$ . Find the values of p and q such that f(1)=3 is an extreme value of f on [0,2]. Is this value a maximum or minimum?
- b) Show that  $y=xe^{\frac{-x^2}{2}}$  satisfies the equation  $xy=(1-x^2)y$
- c) Find the area of the region below the interval [2,1] and above the curve  $y=x^3$
- d) Solve differential equation  $\frac{dy}{dx}y = e^{2x}$ .
- e) Evaluate  $\lim_{(x,y)\to(0,0)} \frac{x^2y^2}{\sqrt{(x^2+y^2)^2}}$  by converting to polar coordinates.

