(Time: 2¹/₂ hours)

Total Marks: 75

N. B.: (1) <u>All</u> questions are <u>compulsory</u>.

- (2) Make suitable assumptions wherever necessary and state the assumptions made.
- (3) Answers to the <u>same question</u> must be <u>written together</u>.
- (4) Numbers to the <u>**right**</u> indicate <u>**marks**</u>.
- (5) Draw <u>neat labeled diagrams</u> wherever <u>necessary</u>.
- (6) Use of **Non-programmable** calculators is **allowed**.

1. Attempt *any three* of the following:

a. Compute the median.

Size	5	6	7	8	9	10	19	12	13
Frequen	48	52	56	60	63	57	55	50	52
су				Â		3.5		2 A 2	N 8 2

- b. Define Geometric Mean. How to calculate it? Find the G.M. of 250, 12, 4.5, 119.5, 42, 35.4, 75, 30
- c. From the following data, calculate Q_1 and Q_3 .

Marks more than	$\langle 10 \rangle$	20	30	40	50	60	70
No. of Students	12	30	54	76	91	101	102

d. Calculate the standard deviation of the heights of 10 students given as,

	LAV - U LV AL C				
Height (in cms.) 161	162 160	163 160	163 164	164	170 164

e. Find the quartile deviation for the following data.

Marks x	5	10	15	20	25	30
No. of Student f	$\sim 2^{\circ}$	3	8	7.4.5	60	849

f. Define Factors and Data Frames in 'R'. How to create them in 'R'.

2. Attempt *any three* of the following:

- a. Explain the Relation between Raw Moments and Central Moments.
- b. Define Skewness. Compute Coefficient of Skewness for the following observations 2, 3, 5, 7, 4, 8, 1.
- c. A random variable X has the following probability distribution values of X

	$\mathbf{X} < 0$		122.2	2 3 3	3	4	5	6	7
5	P(X) 0		k s S	2k	2k	3k	k ²	$2k^2$	$7k^{2} + k$
0		17 - M		110 m (mm.	α (\cdot)		-		

Find (i) k (ii) P(X<6) (iii) P(X≥6) (iv) P(0<X<5)

[TURN OVER]

15

15

- d. The data from a survey of 140 students showed that 37 study Music, 103 play a sport and 25 do neither. Create a Venn diagram to illustrate the data collected and then determine the probability that if a student is selected at random.
 - a) He or she will study music
 - b) He or she will study music given that he or she play sport
- e. It has been found that 2% of the tools produced by a certain machine are defective. What is the probability that in a shipment of 400 such tools.
 - a) 3% or more
 - b) 2% or less

will prove defective?

- f. The electric light bulbs of manufacturer A have a mean lifetime of 2800 hours with a standard deviation of 400 hours. While those of manufacturer B have a mean lifetime of 2400 hours with standard deviation of 200 hours. If random samples of 250 bulbs of each brand are tested, what is the probability that the brand A bulbs will have a mean lifetime that is at least,
 - (a) 320 hours
- (b) 500 hours more than the brand B bulbs?

3. Attempt *any three* of the following:

15

- a. Five bolts drawn From the bolts produced by a certain machine have lengths 3.3, 3.28, 3.31, 3.33, 3.28 cms. Find an unbiased estimate of the variance of the length the bolts produced by the machine
- b. A random sample of 100 articles selected from a batch of 2000 articles shows the average diameter of the articles is 0.354 with standard deviation 0.048. find 95% confidence interval for the average whole batch.
- c. How to calculate an Interval Estimate? State the difference between a Point Estimate and an Interval Estimate.
- d. For a certain test for coin; $H_0: P = \frac{1}{2}$ against $H_1: P = \frac{1}{3}$ can be retained where P represents

the probability of getting a tail. To decide this coin is tossed four times and H_0 reject only if the number of heads observed is 0 or 1. Find the probability of both types of errors.

- e. A certain coin is showed up head 270 occasions in 500 tosses. Test the claim that the coin is unbiased.
- f. Explain with example in-built functions available in 'R' to generate normal distribution.

[TURN OVER]

4. Attempt *any three* of the following:

- a. For a random sample of 10 pigs fed on diet A, the increases in weight in pounds in a certain period were:
 10, 6, 16, 17, 13, 12, 8, 14, 15, 9 lbs.
 For another sample of 12 pigs, fed on diet B, the increases in the same period were.
 7, 13, 22, 15, 12, 14, 18, 8, 21, 23, 10, 17 lbs
 Test whether diets A and B differ significantly in weight. Take α = 0.05
- b. Test the significance of the following rank correlation coefficient at 5% level. R = 0.139, n = 10
- c. Suppose 3 groups each consisting of four students are taken from class XII from 3 different schools. A test of mathematics is taken. The results is:

School A :	71, 75, 65, 69
School B :	90, 80, 86, 84
School C :	72, 77, 76, 79

Compare the mean scores of the schools.

d. In an experiment to study the independence of hypertension on smoking habits, the following data are taken from 180 individuals.

	Non- smokers	Moderate smokers	Heavy smokers	Total
Hypertension	21	36	3000	87
No-hypertension	48	26	19	93
Total	69	62	49	180

Test the hypothesis at 0.05 level of significance that the presence or absence of hypertension is independent of smoking habits.

e. The number of defects per unit in a sample of 330 units of a manufactured product was found as follow:

No. of defects	0	$\sim 1_{\odot}$	>2	\$ 3 3	432
No. of units	2	92	20	3	
	1		1978) 1978)		
0.81.62.62.62.60.66	4	DV NA	1.0° 1×	55	ST ST

Fit a position distribution to the data and test the goodness of fit.

f. How to calculate Contingency Coefficient?

[TURN OVER]

15

5. Attempt *any three* of the following:

a. Find the regression lines of equation for the following.

				62.5	2.0.
Advertising Expenditure ('000 Rs.)	3	5	7	ିହ	212
Quarterly Sales ('0000 units)	9	12	16	14°	15

b. Fit a second degree curve $Y = a+bx+cx^2$ to the following data

X	5	10	15	20	25 30
У	11	13	16	20	28 36

c. Calculate the Coefficient of Correlation between the Age and Blood pressure of given people from a colony.

Age in Years	60	65	70	40	45	50	55
Blood Pressure	145	160	160	125	140	140	145

d. Given the following data estimate the linear trend equation. Find trend values and calculate the trend value of

Year A	2010	2011	2012	2013	2014
No. of cars (in Thousand)	11	30	38	50	56

e. Find (a) σ_x (b) σ_y (c) V(x) (d) V(y) and (e) cov (x, y) for the following data:

X	3
У	
. 1	1100

Also verify $r = \sigma xy / \sigma x . \sigma y$

f. The two regression lines between x and y are given below. Find \overline{x} , \overline{y} and r. 100y - 45x - 1400 = 04y - 5x + 200 = 0